



Wisconsin Department of Agriculture, Trade and Consumer Protection  
 Division of Agricultural Resource Management  
 Bureau of Land and Water Resources  
 PO Box 8911, Madison WI 53708-8911, Phone: 608-224-4605

Use this form to check nutrient management (NM) plans for compliance with the WI NRCS 2015-590 Standard.

# Nutrient Management Checklist Wis. Stat. §92.05(3) (k), Wis. Admin. Code §ATCP50.04(3) and Ch. 51

COUNTY	DATE PLAN SUBMITTED	GROWING SEASON YEAR PLAN IS WRITTEN FOR	(from harvest to harvest)	
TOWNSHIP: (T. N.)	RANGE: (R. E., W.)	CHECK ONE: <input type="checkbox"/> Initial Plan or <input type="checkbox"/> Updated Plan		
NAME OF FARM OPERATOR RECEIVING NM PLAN First Name LastName		FARM NAME (OPTIONAL)		BUSINESS PHONE ( ) -
STREET ADDRESS		CITY	STATE	ZIP
REASON THE PLAN WAS DEVELOPED: <b>Click and choose.</b> (Ordinance, NR 243 WPDES or NOD, DATCP-FP or cost share (cs), DNR-cs, USDA-cs, Other)				CROPLAND ACRES (OWNED & RENTED)
RENTED FARM(S) LANDOWNER NAME(S) AND ACREAGE: add sheet(s) if needed				
WAS THE PLAN WRITTEN IN SNAPPLUS?		<input type="checkbox"/> YES <input type="checkbox"/> NO		If yes, which software version, if known?
CHECK PLANNER'S QUALIFICATION: <b>Click and choose.</b> (1. NAICC-CPCC, 2. ASA-CCA, 3. SSSA-Soil Scientist, 4. DATCP approved training course, 5. Other approved by DATCP)				
NAME OF QUALIFIED NUTRIENT MANAGEMENT PLANNER First Name Last Name			BUSINESS PHONE ( ) -	
STREET ADDRESS		CITY	STATE	ZIP

Use header sections to add comments. Mark NA in the shaded sections if no manure is applied.

1. Does the plan include the following nutrient application requirements to protect surface and groundwater?			
<i>This section applies to fields and pastures. If no manure is applied, check NA for 1.c., 1.h., 1.i., 1.n., 1.o., 1.q., 1.s.</i>	<b>Yes</b>	<b>No</b>	<b>NA</b>
a. Determine field nutrient levels from soil samples analyzed by a DATCP <b>certified laboratory</b> .	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. For fields or pastures with mechanical nutrient applications, determine field nutrient levels from <b>soil samples</b> collected within the last <b>4 years</b> according to 590 Standard (590) and UWEX Pub. A2809, <i>Nutrient Application Guidelines for Field, Vegetable, and Fruit Crops in Wisconsin</i> (A2809) typically collecting <b>1 sample per 5 acres</b> of 10 cores. Soil tests are not required on <b>pastures</b> that do not receive mechanical applications of nutrients if either of the following applies: 1. The pasture average stocking rate is one animal unit per acre or less at all times during the grazing season. 2. The pasture is winter grazed or stocked at an average stocking rate of more than one animal unit per acre during the grazing season, and a nutrient management plan for the pasture complies with 590 using an assumed soil test phosphorus level of 150 PPM and organic matter content of 6%.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. For <b>livestock siting permit approval</b> , collect and analyze soil samples meeting the requirements above in 1. b., excluding pastures, within 12 months of approval and revise the nutrient management plan accordingly. Until then, either option below maybe used: 1. Assume soil test phosphorus levels are greater than 100 ppm soil test P, OR 2. Use <u>preliminary estimates analyzed by a certified DATCP laboratory with soil samples representing &gt; 5 ac/sample.</u>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. <b>Identify all fields'</b> name, boundary, acres, and location.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Use the field's previous year's legume credit and/or applications, predominant soil series, and realistic yield goals to determine the crop's nutrient <b>application rates consistent with A2809 for ALL forms of N, P, and K.</b>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Make no <b>winter applications of N and P</b> fertilizer, except on grass pastures and winter grains.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Document method used to determine <b>application rates</b> . Nutrients shall not runoff during or immediately after application.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Identify in the plan that <b>adequate acreage</b> is available for manure produced and/or applied.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Apply a single phosphorus (P) assessment using either the <b>P Index</b> or <b>soil test P</b> management strategy to all fields within a tract when fields receive manure or organic by-products during the crop rotation.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Use <b>complete crop rotations</b> and the field's <b>critical soil</b> series to determine that sheet and rill erosion estimates will not exceed <b>tolerable soil loss (T)</b> rates on fields that receive nutrients.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Use contours; reduce tillage; adjust the crop rotation; or implement other practices to <b>prevent ephemeral erosion</b> ; and maintain perennial vegetative cover to <b>prevent reoccurring gullies</b> in areas of concentrated flow.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
l. Make no nutrient applications within 8' of <b>irrigation wells</b> or where <b>vegetation is not removed</b> .	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
m. Make no nutrient applications within <b>50' of all direct conduits to groundwater</b> , unless directly deposited by <b>gleaning/pasturing animals</b> or applied as starter fertilizer to corn.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

	Yes	No	NA
n. Make no <b>untreated manure applications</b> to areas within 1000' of a <b>community potable water well</b> or within 100' of a <b>non-community potable water well</b> (ex. church, school, restaurant) unless manure is treated to substantially eliminate pathogens.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
o. Make no manure applications to areas <b>locally delineated</b> by the Land Conservation Committee or in a conservation plan as areas contributing runoff to direct conduits to groundwater unless manure is substantially buried within 24 hours of application.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
p. Make no applications of <b>late summer or fall commercial N fertilizer</b> to the following areas UNLESS needed for establishment of fall seeded crops OR to meet A2809 with a blended commercial fertilizer. Commercial fertilizer N applications shall not exceed 36 lbs. N/acre on: <ul style="list-style-type: none"> <li>• Sites vulnerable to N leaching <b>PRW Soils</b> (P=high permeability, R= bedrock &lt; 20 inches, or W= wet &lt; 12 inches to apparent water table);</li> <li>• Soils with depths of 5 feet or less to bedrock;</li> <li>• Area within 1,000 feet of a community potable water well.</li> </ul> <b>On P soils, when commercial N is applied</b> for full season crops in <b>spring and summer</b> , follow A2809 and apply one of the following: <ol style="list-style-type: none"> <li>1. A split or delayed N application to apply a majority of crop N requirement after crop establishment.</li> <li>2. Use a nitrification inhibitor with ammonium forms of N.</li> <li>3. Use slow and controlled release fertilizers for a majority of the crop N requirement applied near the time of planting.</li> </ol>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
q. Limit manure applications in late summer or fall using the lesser of A2809 or the following 590 rates on <b>PRW Soils</b> . <u>Use ≤ 120 lbs. available N/acre on:</u> <b>P and R soils</b> on <i>all crops, except annual crops</i> . Additionally, manure with ≤ 4% dry matter (DM) wait until after soil temp. < 50°F or Oct. 1, and use either a nitrification inhibitor OR surface apply and do not incorporate for at least 3 days. <b>W soils or combo. W soils</b> on <i>all crops</i> . Additionally, manure with ≤ 4% DM on <i>all crops</i> use at least one of the following: <ol style="list-style-type: none"> <li>1. Use a nitrification inhibitor;</li> <li>2. Apply on an established cover crop, an overwintering annual, or perennial crop;</li> <li>3. Establish a cover crop within 14 days of application;</li> <li>4. Surface apply &amp; don't incorporate for at least 3 days;</li> <li>5. Wait until after soil temp. &lt; 50°F or Oct. 1.</li> </ol> <u>Use ≤ 90 lbs. available N/acre on:</u> <b>P and R soils</b> on <i>annual crops</i> wait until after soil temp. < 50°F or Oct. 1. Additionally, manure with ≤ 4% DM use either a nitrification inhibitor OR surface apply and do not incorporate for at least 3 days. <b>W soils or combination W soils</b> receiving manure with ≤ 4% DM on <i>all crops</i> .	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
r. Use at least one of the following practices on <b>non-frozen soils for all nutrient applications</b> within Surface Water Quality Management Area ( <b>SWQMA</b> ) = 1000' of lakes/ponds or 300' of rivers: <ol style="list-style-type: none"> <li>1. Maintain &gt; 30% cover after nutrient application;</li> <li>2. Effective incorporation within 72 hours of application;</li> <li>3. Establish crops prior to, at, or promptly following application;</li> <li>4. Install/maintain vegetative buffers or filter strips;</li> <li>5. Have at least 3 consecutive years no-till for applications to fields with &lt; 30% residue (silage) and apply nutrients within 7 days of planting.</li> </ol>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
s. Limit mechanical applications to <b>12,000 gals/acre of unincorporated liquid manure or organic by-products</b> with 11% or less dry matter where <b>subsurface drainage</b> is present OR within <b>SWQMA</b> . Wait a minimum of 7 days between sequential applications AND use one or more of the practice options on non-frozen soils listed in 1.r.1. through 1.r.5.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. When frozen or snow-covered soils prevent effective incorporation, does the plan follow these requirements for winter applications of all mechanically applied manure or organic by-products? <i>This section doesn't apply to winter gleaning/pasturing meeting 590 N and P requirements.</i>			
<i>If no manure is applied, check NA for 2.a. through 2.g..</i>			
a. Identify <b>manure quantities planned to be spread during the winter</b> , or the amount of manure generated in 14 days, whichever is greater. <i>For daily haul systems, assume 1/3 of the manure produced annually will need to be winter applied.</i>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Identify <b>manure storage capacity</b> for each type applied and stacking capacity for manure ≥ 16% DM if permanent storage does not exist.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Show on map and make no applications within the <b>SWQMA</b> .	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Show on map and make no surface applications of liquid manure during <b>February and March</b> where <b>Silurian dolomite</b> is within 60 inches of the soils surface OR where <b>DNR Well Compensation</b> funds provided replacement water supplies for wells contaminated with livestock manure.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Show on map and make no applications of manure within <b>300 feet of direct conduits to groundwater</b> .	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Do not exceed the P removal of the following growing season's crop when applying manure. Liquid manure applications are limited to <b>7,000 g/acre</b> . All winter manure applications are not to exceed <b>60 lbs. of P2O5/acre</b> .	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Make no applications of manure to fields with <b>concentrated flow channels</b> unless using two of the following: <ol style="list-style-type: none"> <li>1. Contour buffer strips or contour strip cropping;</li> <li>2. Leave all crop residue and no fall tillage;</li> <li>3. Apply manure in intermittent strips on no more than 50% of field;</li> <li>4. Apply manure on no more than 25% of the field waiting a minimum of 14 days between applications;</li> <li>5. Reduce manure app. rate to 3,500 gal. or 30 lbs. P2O5, whichever is less;</li> <li>6. No manure application within 200 feet of all concentrated flow channels;</li> <li>7. Fall tillage is on the contour and slopes are lower than 6%.</li> </ol> Make no applications to <b>slopes greater than 6%</b> (soil map units with C, D, E, and F slopes) unless the plan documents that no other accessible fields are available for winter spreading AND two of the options 2.g.1. through 2.g.5. are used.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**I certify that the plan represented by the answers on this checklist complies with Wisconsin's NRCS 2015-590 NM Standard or is otherwise noted.**

Qualified NM planner signature	NAICC-Certified Professional Crop Consultant, ASA-Certified Crop Adviser, or SSSA-Soil Scientist	Date
Qualified NM farmer-planner or Authorized farm operator signature receiving and understanding the plan	Date	Signature if reviewed for quality assurance Date